# **Complete Summary**

#### TITLE

Surgical care improvement project: percent of surgical patients with urinary catheter removed on postoperative day 1 or postoperative day 2 with the day of surgery being day zero.

# SOURCE(S)

Specifications manual for national hospital inpatient quality measures, version 3.0c. Centers for Medicare & Medicaid Services (CMS), The Joint Commission; 2009 Oct 1. various p.

# **Measure Domain**

## **PRIMARY MEASURE DOMAIN**

**Process** 

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the Measure Validity page.

## SECONDARY MEASURE DOMAIN

Does not apply to this measure

## **Brief Abstract**

# **DESCRIPTION**

This measure is used to assess the percent of surgical patients with urinary catheter removed on postoperative day 1 (POD 1) or postoperative day 2 (POD 2) with the day of surgery being day zero.

#### **RATIONALE**

It is well–established that the risk of catheter-associated urinary tract infection (UTI) increases with increasing duration of indwelling urinary catheterization. In 2000, Saint reported the results of a pooled analysis of 10 prospective trials dating from 1983 to 1995 which estimated that bacteriuria will develop in 26% of patients after 2 to 10 days of catheterization (95% CI 23-25%). Additional pooled analyses demonstrated that 24% (95% CI 16% to 32%) of those patients will develop symptomatic UTI and bacteremia will develop in 3.6%. Among surgical

patients, two studies of postoperative patients discharged to subacute care with urinary catheters were more likely to be readmitted to the hospital with a UTI compared with those who had catheters removed prior to hospital discharges (Wald, 2005 and Wald, 2008). Among selected major surgical patients in the Surgical Infection Project (SIP) cohort, Wald demonstrated (in press) that 85% had perioperative indwelling catheters placed and half of those patients had catheters for greater than 2 days postoperatively. These patients were twice as likely to develop UTIs prior to hospital discharge. On multivariate analysis, those who had indwelling bladder catheters for more than 2 days postoperatively were 21% more likely to develop UTI, significantly less likely to be discharged to home, and had a significant increase in mortality at 30 days. Additional analyses suggest that there is sizeable variation in the duration of postoperative catheterization among hospitals and that hospital factors may account for this variation. In 2006, Stephan reported the results of a multifaceted intervention study in orthopedic surgery patients in which protocols limiting the use and duration of postoperative catheterization played a large role. They reported a resultant 60% reduction in UTI incidence-density.

## PRIMARY CLINICAL COMPONENT

Surgical care infection prevention; urinary catheter

#### **DENOMINATOR DESCRIPTION**

All selected surgical patients with a catheter in place postoperatively (see the related "Denominator Inclusions/Exclusions" field in the Complete Summary and Appendix A, Table 5.10 for the list of selected surgeries)

#### NUMERATOR DESCRIPTION

Number of surgical patients whose urinary catheter is removed on postoperative day 1 (POD 1) or postoperative day 2 (POD 2) with day of surgery being day zero

# **Evidence Supporting the Measure**

# **EVIDENCE SUPPORTING THE CRITERION OF QUALITY**

 One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

# **Evidence Supporting Need for the Measure**

## **NEED FOR THE MEASURE**

Use of this measure to improve performance

## **EVIDENCE SUPPORTING NEED FOR THE MEASURE**

Saint S. Clinical and economic consequences of nosocomial catheter-related bacteriuria. Am J Infect Control2000 Feb;28(1):68-75. <a href="PubMed">PubMed</a>

Stephan F, Sax H, Wachsmuth M, Hoffmeyer P, Clergue F, Pittet D. Reduction of urinary tract infection and antibiotic use after surgery: a controlled, prospective, before-after intervention study. Clin Infect Dis2006 Jun 1;42(11):1544-51. PubMed

Wald H, Epstein A, Kramer A. Extended use of indwelling urinary catheters in postoperative hip fracture patients. Med Care2005 Oct;43(10):1009-17. PubMed

Wald HL, Epstein AM, Radcliff TA, Kramer AM. Extended use of urinary catheters in older surgical patients: a patient safety problem. Infect Control Hosp Epidemiol2008 Feb;29(2):116-24. <a href="PubMed">PubMed</a>

Wald HL, Ma A, Bratzler DW, Kramer AM. Indwelling urinary catheter use in the postoperative period: analysis of the national surgical infection prevention project data. Arch Surg2008 Jun;143(6):551-7. PubMed

## **State of Use of the Measure**

#### STATE OF USE

Current routine use

## **CURRENT USE**

Accreditation
Collaborative inter-organizational quality improvement
External oversight/Medicaid
External oversight/Medicare
Internal quality improvement

# **Application of Measure in its Current Use**

## **CARE SETTING**

Hospitals

## PROFESSIONALS RESPONSIBLE FOR HEALTH CARE

Measure is not provider specific

## **LOWEST LEVEL OF HEALTH CARE DELIVERY ADDRESSED**

Single Health Care Delivery Organizations

## **TARGET POPULATION AGE**

Age greater than or equal to 18 years

# **TARGET POPULATION GENDER**

# STRATIFICATION BY VULNERABLE POPULATIONS

Unspecified

# **Characteristics of the Primary Clinical Component**

## INCIDENCE/PREVALENCE

See the "Rationale" field.

## **ASSOCIATION WITH VULNERABLE POPULATIONS**

Unspecified

# **BURDEN OF ILLNESS**

See the "Rationale" field.

## **UTILIZATION**

Studies of postoperative patients discharged to subacute care with urinary catheters were more likely to be readmitted to the hospital with a urinary tract infection (UTI) compared with those who had catheters removed prior to hospital discharges.

See also the "Rationale" field.

## **EVIDENCE FOR UTILIZATION**

Wald H, Epstein A, Kramer A. Extended use of indwelling urinary catheters in postoperative hip fracture patients. Med Care2005 Oct;43(10):1009-17. PubMed

Wald HL, Epstein AM, Radcliff TA, Kramer AM. Extended use of urinary catheters in older surgical patients: a patient safety problem. Infect Control Hosp Epidemiol2008 Feb;29(2):116-24. PubMed

# **COSTS**

Unspecified

**Institute of Medicine National Healthcare Quality Report Categories** 

## **IOM CARE NEED**

Staying Healthy

## **IOM DOMAIN**

## **Data Collection for the Measure**

#### **CASE FINDING**

Users of care only

## **DESCRIPTION OF CASE FINDING**

All selected surgical patients with a urinary catheter in place postoperatively

## **DENOMINATOR SAMPLING FRAME**

Patients associated with provider

# **DENOMINATOR INCLUSIONS/EXCLUSIONS**

# **Inclusions**

All selected surgical patients with an International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) Principal Procedure Code of selected surgeries as defined in the appendices of the original measure documentation with a catheter in place postoperatively

## **Exclusions**

- Patients less than 18 years of age
- Patients who have a Length of Stay (LOS) greater than 120 days
- Patients who had a principal diagnosis suggestive of preoperative infectious diseases (refer to Appendix A, Table 5.09 of the original measure documentation for ICD-9-CM codes)
- Patients whose ICD-9-CM principal procedure was performed entirely by Laparoscope
- Patients enrolled in clinical trialsÂ
- Patients who had a urological, gynecological or perineal procedure performed (refer to Appendix A, Table 5.16 of the original measure documentation for ICD-9-CM codes)
- Patients whose ICD-9-CM principal procedure occurred prior to the date of admission
- Patients who had other procedures requiring general or spinal anesthesia that occurred within 3 days (4 days for coronary artery bypass graft [CABG] or Other Cardiac Surgery) prior to or after the procedure of interest (during separate surgical episodes) during this hospital stay
- Patients with physician/advanced practice nurse/physician assistant (physician/APN/PA) documented infection prior to surgical procedure of interest
- Patients who expired perioperatively
- Patients whose length of stay was less than two days postoperatively
- Patients who had a suprapubic catheter or had intermittent catheterization preoperatively

- Patients who did not have a catheter in place postoperatively
- Patients who had physician/APN/PA documentation of a reason for not removing the urinary catheter postoperatively

# **RELATIONSHIP OF DENOMINATOR TO NUMERATOR**

All cases in the denominator are equally eligible to appear in the numerator

# **DENOMINATOR (INDEX) EVENT**

Institutionalization
Therapeutic Intervention

#### **DENOMINATOR TIME WINDOW**

Time window brackets index event

# **NUMERATOR INCLUSIONS/EXCLUSIONS**

## **Inclusions**

Number of surgical patients whose urinary catheter is removed on postoperative day 1 (POD 1) or postoperative day 2 (POD 2) with day of surgery being day zero

## **Exclusions**

None

# MEASURE RESULTS UNDER CONTROL OF HEALTH CARE PROFESSIONALS, ORGANIZATIONS AND/OR POLICYMAKERS

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

## **NUMERATOR TIME WINDOW**

Fixed time period

#### **DATA SOURCE**

Administrative data Medical record

# LEVEL OF DETERMINATION OF QUALITY

Individual Case

# PRE-EXISTING INSTRUMENT USED

Unspecified

# **Computation of the Measure**

## **SCORING**

Rate

#### **INTERPRETATION OF SCORE**

Better quality is associated with a higher score

# **ALLOWANCE FOR PATIENT FACTORS**

Unspecified

# STANDARD OF COMPARISON

External comparison at a point in time External comparison of time trends Internal time comparison

# **Evaluation of Measure Properties**

# **EXTENT OF MEASURE TESTING**

Unspecified

# **Identifying Information**

# **ORIGINAL TITLE**

SCIP-Inf-9: urinary catheter removed on postoperative day 1 (POD 1) or postoperative day 2 (POD 2) with day of surgery being day zero.

# **MEASURE COLLECTION**

National Hospital Inpatient Quality Measures

# **MEASURE SET NAME**

Surgical Care Improvement Project (SCIP)

# **SUBMITTER**

Centers for Medicare & Medicaid Services Joint Commission, The

## **DEVELOPER**

# **FUNDING SOURCE(S)**

All external funding for measure development has been received and used in full compliance with The Joint Commission's Corporate Sponsorship policies, which are available upon written request to The Joint Commission.

# **COMPOSITION OF THE GROUP THAT DEVELOPED THE MEASURE**

The Centers for Medicare & Medicaid Services assembled and maintained the Technical Expert Panel for development of the Surgical Infection Prevention Project (SIP) measures in 2002. The SIP set subsequently transitioned to the Surgical Care Improvement Project (SCIP) effective July 1, 2006. The panel has been maintained by the Centers for Medicare & Medicaid Services since the inception of the project.

SCIP Partners include the Steering Committee of 10 national organizations who have pledged their commitment and full support for SCIP:

- Agency for Healthcare Research and Quality
- American College of Surgeons
- American Hospital Association
- American Society of Anesthesiologists
- Association of Perioperative Registered Nurses
- Centers for Disease Control and Prevention
- Centers for Medicare & Medicaid Services
- Institute for Healthcare Improvement
- The Joint Commission
- Veterans Health Administration

# FINANCIAL DISCLOSURES/OTHER POTENTIAL CONFLICTS OF INTEREST

Expert panel members have made full disclosure of relevant financial and conflict of interest information in accordance with the Conflict of Interest policies, copies of which are available upon written request to The Joint Commission and the Centers for Medicare & Medicaid Services.

# **ENDORSER**

National Quality Forum

## **ADAPTATION**

Measure was not adapted from another source.

# **RELEASE DATE**

2009 Oct

## **MEASURE STATUS**

This is the current release of the measure.

# SOURCE(S)

Specifications manual for national hospital inpatient quality measures, version 3.0c. Centers for Medicare & Medicaid Services (CMS), The Joint Commission; 2009 Oct 1. various p.

# **MEASURE AVAILABILITY**

The individual measure, "SCIP-Inf-9: Urinary Catheter Removed on Postoperative Day 1 (POD 1) or Postoperative Day 2 (POD 2) with Day of Surgery Being Day Zero," is published in "Specifications Manual for National Hospital Inpatient Quality Measures." This document is available from <a href="The Joint Commission Web site">The Joint Commission Web site</a>. Information is also available from the <a href="Centers for Medicare & Medicaid Services">Centers for Medicare & Medicaid Services</a> (CMS) Web site. Check The Joint Commission Web site and CMS Web site regularly for the most recent version of the specifications manual and for the applicable dates of discharge.

## **COMPANION DOCUMENTS**

The following are available:

- A software application designed for the collection and analysis of quality improvement data, the CMS Abstraction and Reporting Tool (CART), is available from the <u>CMS CART Web site</u>. Supporting documentation is also available. For more information, e-mail CMS PROINQUIRIES at proinquiries@cms.hhs.gov.
- The Joint Commission. A comprehensive review of development and testing for national implementation of hospital core measures. Oakbrook Terrace (IL): The Joint Commission; 40 p. This document is available from <u>The Joint Commission Web site</u>.
- The Joint Commission. Attributes of core performance measures and associated evaluation criteria. Oakbrook Terrace (IL): The Joint Commission;
   5 p. This document is available from The Joint Commission Web site.

# **NQMC STATUS**

The Joint Commission submitted this NQMC measure summary to ECRI Institute on July 31, 2009. This NQMC summary was reviewed accordingly by ECRI Institute on December 2, 2009. The information was verified by the Centers for Medicare & Medicaid Services on February 18, 2010.

#### **COPYRIGHT STATEMENT**

The Specifications Manual for National Hospital Inpatient Quality Measures [Version 3.0b, October, 2009] is the collaborative work of the Centers for Medicare & Medicaid Services and The Joint Commission. The Specifications Manual is periodically updated by the Centers for Medicare & Medicaid Services and The Joint Commission. Users of the Specifications Manual for National

Hospital Quality Measures should periodically verify that the most up-to-date version is being utilized.

#### Disclaimer

# **NQMC DISCLAIMER**

The National Quality Measures Clearinghouse™ (NQMC) does not develop, produce, approve, or endorse the measures represented on this site.

All measures summarized by NQMC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public and private organizations, other government agencies, health care organizations or plans, individuals, and similar entities.

Measures represented on the NQMC Web site are submitted by measure developers, and are screened solely to determine that they meet the NQMC Inclusion Criteria which may be found at

http://www.qualitymeasures.ahrq.gov/about/inclusion.aspx.

NQMC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or its reliability and/or validity of the quality measures and related materials represented on this site. The inclusion or hosting of measures in NQMC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding measure content are directed to contact the measure developer.

Copyright/Permission Requests

Date Modified: 4/5/2010

